

State of The Art Management of POSTTRANSPLANT SEQUELAE

FOR PHYSICIANS,
NURSES, AND PHARMACISTS

PRESENTED BY:



NATIONAL INSTITUTE OF ALLERGY AND INFECTIOUS DISEASES OF
THE NATIONAL INSTITUTES OF HEALTH
U S DEPARTMENT OF HEALTH AND HUMAN SERVICES

Impact and Management of Cardiovascular Risks After Kidney Transplantation

FIRST IN A SERIES OF MONOGRAPHS

BASED ON A ROUNDTABLE

HELD JANUARY 28, 2005,

IN BETHESDA, MARYLAND

THIS PROGRAM IS SUPPORTED BY
AN UNRESTRICTED EDUCATIONAL
GRANT FROM WYETH.

IN COOPERATION WITH:



AMERICAN SOCIETY OF
TRANSPLANT SURGEONS



UNITED NETWORK
FOR ORGAN SHARING



NORTH AMERICAN TRANSPLANT
COORDINATORS ORGANIZATION



INTERNATIONAL TRANSPLANT
NURSES SOCIETY

JOINTLY SPONSORED BY:

UNIVERSITY
OF MINNESOTA



Accreditation

This activity has been planned and implemented in accordance with the Essential Areas and Policies of the Accreditation Council for Continuing Medical Education (ACCME) and the Criteria for Quality of the Accreditation Council for Pharmacy Education (ACPE) through the joint sponsorship of the University of Minnesota Office of Continuing Medical Education, University of Minnesota Office of Continuing Pharmacy Education, and SynerMed® Communications.

CME-Accredited Sponsor

The University of Minnesota is accredited by the ACCME to provide continuing medical education for physicians (Provider Number 0000810).

CPE Accreditation



The University of Minnesota College of Pharmacy is accredited by the ACPE as a provider of continuing pharmacy education. The Universal Program Number is 031-999-05-073-H01. Following completion of the program, successful passing of the posttest (70% or better), and submission of a program evaluation, statements of credit will be mailed to participants within 4 weeks of receipt of materials.

Credit Designation Statement

Physicians—The University of Minnesota designates this educational activity for a maximum of 1.0 category 1 credit toward the AMA Physician's Recognition Award. Each physician should claim only those credits that he/she actually spent in the educational activity.

Nurses—This program was designed to meet Minnesota Board of Nursing continuing education requirements and provides 1.2 contact hours of continuing education.

Pharmacists—The University of Minnesota approves this educational activity for 1.0 contact hour of pharmacy continuing education credit.

Financial Support

This program is supported by an unrestricted educational grant from Wyeth.

Educational Objectives

At the conclusion of this program, participants will be able to:

- Recognize the prevalence and long-term impact of overall cardiovascular disease, and specifically hypertension and dyslipidemias, in renal transplant recipients
- Describe the risk factors for hypertension and dyslipidemias
- Identify the effects of immunosuppressive drugs on the development of hypertension and dyslipidemias
- Outline the modifications to immunosuppressive regimens to help manage hypertension and dyslipidemias
- Discuss the interventions to treat hypertension and dyslipidemias

Target Audience

Transplant surgeons, transplant nephrologists, transplant nurses, transplant coordinators, pharmacists, and other healthcare professionals who are involved in the treatment and management of renal transplant recipients

Term of Approval

Date of Original Release: June 2005

Expiration Date: June 30, 2006

Faculty Disclosure Information

The University of Minnesota endorses the standards of the Accreditation Council for Continuing Medical Education, the Accreditation Council for Pharmacy Education, and the guidelines of the Association of American Medical Colleges that the sponsors of continuing medical education activities and the speakers at these activities disclose significant relationships with commercial companies whose products or services are discussed in educational presentations. For speakers, significant relationships include receiving from a commercial company research grants, consultancies, honoraria and travel, or other benefits or having a self-managed equity interest in a company. Disclosure of a relationship is not intended to suggest or condone bias in any presentation, but is made to provide participants with information that might be of potential importance to their evaluation of a presentation.

Affiliation/Financial Interest			
Faculty Member	Grants/Research Support	Consultant	Speakers Bureau
J. J. Curtis, MD	No affiliation or financial interest		
O. Gaber, MD	Fujisawa Healthcare, Inc; Genzyme Corporation; Novartis; Roche Laboratories Inc; Wyeth-Ayerst Laboratories		Wyeth-Ayerst Laboratories
B. L. Kasiske, MD	Bristol-Myers Squibb; Merck & Co, Inc; Pfizer Inc		Amgen; Fujisawa Healthcare Inc; Merck & Co, Inc; Novartis; Pfizer Inc; Roche Laboratories Inc; Wyeth-Ayerst Laboratories
R. H. Rubin, MD		Merck & Co, Inc; Merrimack; Pfizer Inc; Roche Laboratories Inc	
R. P. Winsett, PhD	No affiliation or financial interest		
E. S. Woodle, MD	Enzon, Inc; Fujisawa Healthcare, Inc; Genzyme Corporation; Roche Laboratories Inc; Wyeth-Ayerst Laboratories		

Impact and Management of Cardiovascular Risks After Kidney Transplantation

Presented by

National Institute of Allergy and Infectious Diseases
National Institutes of Health
Division of Allergy, Immunology, and Transplantation
Bethesda, Maryland

Represented by

Shiv A. Prasad, PhD

Chief, Transplantation Immunobiology Branch
National Institute of Allergy and Infectious Diseases
National Institutes of Health
Bethesda, Maryland

Program Chair

Bertram L. Kasiske, MD

Professor of Medicine
University of Minnesota
Minneapolis, Minnesota

Faculty

John J. Curtis, MD

Professor of Transplant Nephrology
Professor of Surgery, Division of Transplantation
University of Alabama at Birmingham
Birmingham, Alabama

Osama Gaber, MD

Professor, Department of Surgery
Associate Professor, College of Nursing
University of Tennessee Health Science Center
Memphis, Tennessee

Robert H. Rubin, MD

Osborne Professor of Health Sciences and Technology
Professor of Medicine, Harvard Medical School
Associate Director, Division of Infectious Disease
Brigham and Women's Hospital
Director, Center for Experimental Pharmacology and Therapeutics,
Harvard University-Massachusetts Institute of Technology
Division of Health Sciences and Technology
Boston, Massachusetts

E. Steve Woodle, MD

Professor of Surgery
Director, Division of Transplantation
Chairman, Board of Directors
Israel Penn International Transplant Tumor Registry
Department of Surgery
University of Cincinnati College of Medicine
Cincinnati, Ohio

Rebecca P. Winsett, PhD

Associate Professor, College of Nursing
University of Tennessee Health Science Center
Memphis, Tennessee

TABLE OF CONTENTS

PAGE


Introduction	1
Hypertension	1
Estimates of the Prevalence of Hypertension and Its Relationship to Outcomes	1
Factors Associated With Posttransplant Hypertension	2
Management of Hypertension.	3
Diuretics	3
Angiotensin-Converting Enzyme (ACE) Inhibitors and Angiotensin II Receptor Blockers (ARBs)	3
Calcium Channel Blockers (CCBs)	4
ACE Inhibitors Versus CCBs	4
Dyslipidemias	4
The Prevalence of Dyslipidemias and Their Relationship to Morbidity and Mortality.	4
Factors Associated With Posttransplant Dyslipidemias	5
Management of Dyslipidemias	5
The Role of Immunosuppressive Drugs in Dyslipidemias.	6
Corticosteroids	6
Calcineurin and Target of Rapamycin Inhibitors.	6
Role of Lipid-Lowering Drugs	7
Conclusions	7
References	8
Posttest	9
Evaluation	10

INTRODUCTION

Remarkable progress over the last decades due to advances in immunosuppressive therapies and transplantation technology has resulted in improved 1-year kidney graft survival, now exceeding 90%.¹ Although transplantation for patients with end-stage renal disease (ESRD) confers a survival advantage superior to that of maintenance dialysis,² longer-term morbidity and mortality for the transplant recipient are closely associated with development of cardiovascular disease. Currently, cardiovascular disease accounts for nearly 40% of deaths among renal transplant recipients (Figure 1), a rate that is considerably greater than that for the general population and more dramatic for younger than for older patients (Figure 2).^{3,4}

Reviews of available evidence conclude that kidney transplant recipients have many of the same risk factors for cardiovascular disease as those traditionally found in the general population, such as risk factors recognized in the Framingham Heart Study (FHS)^{5,6} and in the National Cholesterol Education Program (NCEP). However, additional cardiovascular disease risk factors have been identified for transplant recipients. Although an observational study found that the risk calculated from the FHS did predict ischemic events in kidney transplant recipients at more than 1 year posttransplantation, the risks were underestimated for the transplant population compared to the general population. Specifically, the increased risks among transplant recipients were associated with diabetes mellitus and, to a lesser degree, age and smoking.⁵ Risk factors unique to transplant recipients that also may


Figure 2 Annual Cardiovascular Mortality for Renal Transplant Recipients and the General Population by Age Group³



Adapted with permission from Foley RN, et al. *Am J Kidney Dis.* 1998;32:S112-S119.

be more critical than the traditional risk factors include decreased renal function, development of proteinuria, and use of immunosuppressive medications.⁷⁻¹⁰ Although lipid abnormalities and hypertension are common risk factors for both transplant recipients and the general population,⁵ the very high incidence of these factors among kidney transplant recipients compels further investigation of their prevalence and potential causes, as well as identification of modifiable factors and therapeutic interventions. The National Kidney Foundation offers guidelines for the management of dyslipidemias and hypertension.¹¹⁻¹³

Figure 1 Causes of Mortality in Renal Transplant Recipients¹⁴



USRDS = United States Renal Data System; CVD = cardiovascular disease.
Courtesy of Bertram L. Kasiske, MD


HYPERTENSION

Estimates of the prevalence of hypertension and its relationship to outcomes

Hypertension in the kidney transplant recipient is extremely common, and, despite the variety of antihypertensive medications available, management of hypertension remains a challenge. A recent analysis portrays the situation now encountered in the transplant community with estimates of 87.6% of patients with blood pressure (BP) higher than normal (normal systolic BP defined as <120 mm Hg) and only 3.5% with truly normal BP in the absence of antihypertensive medication at 1 year posttransplantation.¹⁵ Even this prevalence may be an underestimate since ambulatory BP monitoring of kidney transplant recipients revealed that office BP measurements fail to detect hypertension in 15% of patients considered normotensive.¹⁶

Hypertension is an independent risk factor for poorer long-term kidney graft survival in a variety of different studies.^{15,17,18} Graft and patient survival rates analyzed over 7 years from data collected through the Collaborative Transplant Study international registry showed that increased levels of systolic and diastolic BP 1 year after transplantation were associated with gradual increases in risk of graft failure ($P<.0001$ at 7 years) (Figure 3).¹⁷ A report from the United States further showed that each 10 mm Hg increase in systolic BP was associated with an increased risk of graft failure ($P<.0001$), death-censored graft failure ($P<.0001$), and death ($P<.0001$), independent of acute rejection and other transplant variables. Some of the relationship of BP to graft failure was evident at least 5 years prior to graft loss, but at 1 year posttransplantation 45.5% of patients had uncontrolled hypertension (systolic BP ≥ 140 mm Hg).¹⁵

Figure 3
Association of Systolic Blood Pressure (SBP) With Decreased Graft Survival¹⁷



Adapted with permission from Opelz G, et al. *Kidney Int.* 1998;53:217-222.

Factors associated with posttransplant hypertension

Some causes of posttransplant hypertension include poor allograft function,¹⁹ source of the donor kidney,⁸ retention of the native diseased kidneys,^{15,20} transplant renal artery stenosis,²¹ and immunosuppressive drugs.

The origin of essential hypertension is a practical issue for the transplant community in deciding whether organs from hypertensive donors should be accepted or if more emphasis should be placed on other mechanisms of BP control.²² Whereas some investigators believe that

essential hypertension is not caused by the donor kidney but is more of a systemic disease that can recur,⁸ the physiologist Guyton postulated that the kidney is the prime factor in hypertension.²³ The central role of the kidney was illustrated in a pivotal study of patients with essential hypertension who required kidney transplantation. The recipients remained normotensive for 4.5 years after receiving a kidney from a normotensive donor, with BPs not significantly different from those of the normotensive control subjects.²⁴ Overall, there is consensus that essential hypertension is kidney disease.

Use of some immunosuppressive drugs has become associated with subsequent hypertension. Corticosteroids are associated with only a 15% incidence of hypertension, and corticosteroid withdrawal does have a positive impact on BP control.^{10,25} Paradoxically, although the introduction of calcineurin inhibitors (CNIs) benefited the transplant recipient with greatly increased graft survival, with their widespread use came an increased incidence of hypertension.⁷ The first widely used CNI, cyclosporine, increases BP in solid organ transplant recipients with or without concomitant use of corticosteroids and in a dose-related manner. In a retrospective study, the use of cyclosporine for at least 1 year posttransplantation and the use of higher doses of prednisone were associated with higher BP.¹⁵ Hypertension alleviation is possible if cyclosporine is discontinued early.⁷ Since the CNI tacrolimus also produces hypertension,^{10,26} the underlying intracellular mechanism believed responsible for both immunosuppressive activity and elevated BP is the inhibition of calcineurin.²⁷ In contrast, the immunosuppressive agent sirolimus does not appear to cause hypertension because it does not interact with calcineurin.²⁸

In general, evidence suggests that of the CNIs, tacrolimus causes less hypertension than does cyclosporine.²⁸ Two prospective clinical trials in kidney transplantation compared the outcomes of patients randomized to receive either tacrolimus- or cyclosporine-based immunosuppression. The first report showed that after 6 months of follow-up in 557 patients, a significantly lower proportion of patients in the tacrolimus arm had hypertension than in the cyclosporine arm (15.7% vs 23.2%, respectively; $P=.032$).²⁹ In a separate study of 94 patients, the incidence of hypertension was similar in the two groups until the 3-year measurement, at which time a significantly higher percentage of patients were on antihypertensive medication in the cyclosporine group than in the tacrolimus group (74% vs 50%, respectively; $P<.05$).²⁶ The clinical comparison studies of hypertension caused by these two agents are difficult to interpret because if the hypertensive mechanism relies on calcineurin inhibition, the two CNIs should

have similar effects. However, if this is not the observed clinical effect, the question becomes whether CNIs cause hypertension through mechanisms that are independent of their effects on calcineurin.

A multicenter, randomized trial of tacrolimus in combination with either sirolimus or mycophenolate mofetil (MMF) evaluated short-term results at 6 months among 361 kidney transplant recipients. Although patient and graft survival and systolic BP were similar, diastolic BP was higher in the tacrolimus/sirolimus group than in the tacrolimus/MMF group ($P=.02$).³⁰

A viable option for hypertension reduction is to institute early immunosuppressive agent withdrawal or minimization. In a long-term study of sirolimus after early cyclosporine withdrawal, 430 eligible recipients were randomized to remain on cyclosporine/sirolimus/corticosteroid maintenance immunosuppression or to undergo cyclosporine withdrawal with sirolimus/corticosteroid maintenance. After 2 years of follow-up, graft and patient survival and acute rejection rates were similar, but systolic BP was significantly lower in recipients who had only sirolimus/corticosteroid-based immunosuppression than in those continuing the cyclosporine regimen (134 mm Hg vs 141 mm Hg, respectively; $P<.001$).³¹ In contrast, a large, randomized, multicenter European trial (N=833) instituted a controlled withdrawal of the two adjunctive agents, corticosteroids or MMF, from a tacrolimus-based regimen in renal transplant recipients; however, there was no significant difference in rates of patient or graft survival, acute rejection rates, or incidence of hypertension after 6 months.³²

The goal in choosing an immunosuppressive regimen is, of course, to maintain the survival of the patient and graft while reducing long-term side effects. Hypertension is only one of the many potential side effects to consider in the choice of immunosuppressive drugs (Table 1).

Management of Hypertension

Although no clinical trials have prospectively investigated the degree of BP control and long-term outcomes in kidney transplantation, retrospective analyses have indicated that patients with lower BP have better overall survival and graft survival. So antihypertensive drugs are now routinely prescribed for hypertensive kidney transplant recipients, but management may be complicated by the presence of diabetes or proteinuria from poor allograft function. Target BP levels in renal transplant recipients have been recommended by the National Kidney Foundation Kidney Disease Outcomes Quality Initiative (K/DOQI) and the European Best Practice Guidelines Expert Group on Renal Transplantation (Table 2).^{33,34}

Table 1
Side Effect Profiles of Immunosuppressive Agents

	CsA	Tac	Srl	Ster	MMF
Hypertension	++	+	Ø	++	Ø
Hyperglycemia	+	++	Ø	+++	Ø
Renal insufficiency	++	++	Ø	Ø	Ø
Hyperlipidemia	++	+	++	++	Ø
Hyperkalemia	+++	+++	Ø	Ø	Ø
Tremor	Ø	+	Ø	Ø	Ø
Hirsutism	+	Ø	Ø	Ø	Ø
Gingival hyperplasia	+	Ø	Ø	Ø	Ø
Hypophosphatemia	++	++	+	Ø	Ø
Osteoporosis	±	±	Ø	+++	Ø
Malignancy	+	+	?	Ø	+

CsA = cyclosporine; Tac = tacrolimus; Srl = sirolimus; ster = steroids; MMF = mycophenolate mofetil; +++ = severe; ++ = moderate; + = mild; ± = opposite; Ø = none; ? = unknown.

Adapted from Dr Martin Zand.

Diuretics

Although no comparative trials are available for evaluation of diuretics in posttransplant hypertension, there is general agreement that especially in transplant recipients with volume overload, diuretics are necessary for BP control.^{1,10}

Angiotensin-converting enzyme (ACE) inhibitors and angiotensin II receptor blockers (ARBs)

Initial reluctance to use ACE inhibitors after kidney transplantation was originally based on fears of hyperkalemia and abrupt declines in glomerular filtration rate (GFR) in patients with transplant renal artery stenosis.^{1,10,35} However, more recent studies have reinforced the safety and effectiveness of these agents, which also now appear to have genuine application in hypertensive

Table 2
Blood Pressure Target Levels in Renal Transplant Recipients^{33,34}

National Kidney Foundation (K/DOQI Guidelines)	<130/80 mm Hg
EBPG Expert Group on Renal Transplantation	
Without proteinuria	<130/85 mm Hg
With proteinuria	<125/75 mm Hg

K/DOQI = Kidney Disease Outcomes Quality Initiative; EBPG = European Best Practice Guidelines.

transplant recipients with left ventricular hypertrophy or chronic allograft nephropathy,¹⁰ as well as use in the early posttransplant period (first 90 days).³⁶ ACE inhibitors are also effective in reducing both hematocrit and hemoglobin levels in those patients with posttransplant erythrocytosis.^{1,37}

The ACE inhibitors and ARBs may help preserve renal function as well as lower BP.^{38,39} In an early study of 22 kidney transplant recipients, treatment with enalapril reduced proteinuria without changing creatinine clearance.⁴⁰ Furthermore, a Canadian chart review showed that of 177 renal transplant recipients prescribed either an ACE inhibitor or an ARB, mean arterial pressure decreased at each measurement point and significantly from a mean of 92 mm Hg to 86 mm Hg over 3 years ($P<.05$) without a change in creatinine clearance.³⁷ Additional studies have shown renal benefits during treatment with losartan,⁴¹ captopril,⁴¹ and lisinopril.⁴²

Calcium channel blockers (CCBs)

CCBs, particularly the dihydropyridine CCBs, are attractive antihypertensive agents in transplant recipients because they dilate the afferent arteriole, thus reversing CNi-related vasoconstriction that may lead to progressive renal function decline.⁴³ Among many trials conducted in renal transplantation, a multicenter, placebo-controlled study randomized 210 kidney transplant recipients to receive either the CCB isradipine or placebo. At both 3 and 12 months posttransplantation, renal function was significantly better in the treatment group than in the placebo group ($P=.002$ at 3 months and $P=.021$ at 12 months), without a negative impact on the incidence or severity of delayed graft function or acute rejections.⁴⁴ Other CCBs, including amlodipine,⁴⁵ nifedipine,⁴⁶ and felodipine,⁴⁷ have specifically shown benefits for graft functioning in transplant recipients. Therefore, a number of studies have confirmed that dihydropyridine CCBs are effective in improving renal blood flow and GFR.^{1,43} Unfortunately, nondihydropyridine CCBs (eg, verapamil and diltiazem) interact with the cytochrome P450 (CYP) 3A4 isoenzyme, so coadministration with a CNi can lead to noteworthy pharmacokinetic and pharmacodynamic interactions.¹ Additionally, concerns expressed in the nontransplant literature over potential harm of dihydropyridine CCB use in patients with kidney disease⁴⁸ have led to cautionary use in transplantation.

ACE inhibitors versus CCBs

Currently, no single antihypertensive agent or class is recommended over another, but the impression is that most transplant programs use ACE inhibitors first. Studies have been conducted to evaluate any differential benefit of ACE inhibitors and CCBs. A prospective study randomized 154 renal transplant recipients with


hypertension to receive lisinopril or controlled-release nifedipine. In this study, both agents achieved good BP control, but statistically significant increases in GFR from baseline values were observed only in the nifedipine group at 1 year ($P<.0001$). Serum creatinine levels were also significantly decreased in the nifedipine group compared to baseline at 1 year ($P=.013$) but not in the lisinopril group.⁴⁹ A recent large retrospective study evaluated the relationship between BP and antihypertensive use in 1662 renal transplant recipients. Among the antihypertensive drug classes evaluated, only CCBs were associated with a reduction in graft loss (relative risk [RR], 0.81; 95% confidence interval [CI], 0.68-0.96; $P=.0236$), whereas only ACE inhibitors/ARBs were associated with significantly lowering urinary protein excretion ($P=.041$).³⁹ In summary, no single class of antihypertensive agent has been demonstrated to be superior in maintaining renal allograft function. However, the reality is that most kidney transplant recipients will eventually require agents from most antihypertensive classes for BP control.

DYSLIPIDEMIAS

The prevalence of dyslipidemias and their relationship to morbidity and mortality

Dyslipidemias are prevalent in kidney transplant recipients. Among numerous reports monitoring lipid levels after transplantation, two representative studies reported high levels of low-density lipoprotein cholesterol (LDL-C) and low levels of high-density lipoprotein cholesterol (HDL-C) in kidney transplant recipients (Figure 4).^{13,50,51} Cardiovascular disease mortality is increased for transplant recipients compared to the general population

Figure 4
Prevalence of Dyslipidemias in Renal Transplant Recipients Receiving Immunosuppressive Therapy^{13,50,51}



TC = total cholesterol; LDL-C = low-density lipoprotein cholesterol; HDL-C = high-density lipoprotein cholesterol.

(Figure 2), and several observational studies in kidney transplantation have reported a positive relationship between dyslipidemias and cardiovascular disease.¹³

A number of studies have shown correlations between lipid levels and subsequent chronic allograft nephropathy, graft loss, and decreased patient survival. In particular, a large analysis from Europe evaluated the effects of serum cholesterol as a continuous variable on long-term outcomes in renal transplantation. In this retrospective study of 676 recipients receiving primarily azathioprine- or cyclosporine-based immunosuppression and corticosteroids, serum cholesterol levels at 1 year posttransplantation proved an independent predictor of graft failure (death-censored RR, 1.33; 95% CI, 1.15-1.55; $P=.0001$) and patient survival (RR, 1.5; 95% CI, 1.1- 2.1; $P=.01$).⁵² Furthermore, an analysis of 706 renal transplant recipients in the United States revealed that serum triglyceride levels were a risk factor for chronic rejection independent of acute rejection episodes (RR, 1.09; 95% CI, 1.03-1.16 for each 100 mg/dL; $P=.034$).⁵³

Factors associated with posttransplant dyslipidemias

Transplant recipients have many of the same risk factors for dyslipidemias as does the general population, such as obesity, diet, genetic disposition, and insulin resistance. However, nontraditional risk factors, immunosuppressive drugs in particular, may also adversely affect the relationship between lipid levels and cardiovascular disease. Risk factors assessed in the FHS⁶ and the Multiple Risk Factor Intervention Trial⁵⁴ reported an increased risk of death from coronary heart disease with increasing cholesterol levels in the general population, largely because of the negative impact of higher levels of LDL-C. Subsequent trials and a meta-analysis found that reducing total cholesterol and LDL-C levels also reduced coronary events and mortality in the general population and that the reduction in cardiac events and mortality was proportional to the extent of LDL-C reduction.⁵⁵ Risk factors for ischemic heart disease, identified by the FHS, were compared in a single-center study (N=1124) to posttransplant (>1 year) risk factors. A similar relationship between abnormal cholesterol levels and risk for ischemic events for both male and female transplant recipients was found. Even total cholesterol levels of 200 to 239 mg/dL seemed related to increased incidence of transplant ischemic heart disease.⁵

Management of dyslipidemias


The National Kidney Foundation K/DOQI guidelines place transplant recipients in the highest risk

category (per the NCEP Adult Treatment Panel III), along with patients with chronic kidney disease. Accordingly, intervention is recommended when LDL-C levels are 100 mg/dL or higher. The prevailing opinion in the transplant community is that any transplant recipient with elevated LDL-C concentrations should be managed with both drugs and diet. Those patients with exceptionally high triglyceride levels require a slightly different approach (Figure 5).¹³

Few clinical trials have focused on therapeutic lifestyle changes (TLC) for kidney transplant recipients; however, studies in the general population suggest that exercise and weight reduction produce small improvements in cholesterol levels. Consultation with a renal dietician is important to ensure good nutrition, with emphasis on low-fat foods, soluble fiber, and use of plant sterols for fat replacement strategies. Alcohol in moderation should be discussed, but smoking cessation should be actively pursued.¹³ For patients with modest elevations in total cholesterol values (eg, LDL-C 100-129 mg/dL), TLC can be tried for approximately 3 months before adding a lipid-lowering drug. If LDL-C levels are 130 mg/dL or greater, consideration should be given to starting a lipid-lowering agent along with TLC. For most patients, the agent of choice is a statin. However, the statin dose should be reduced in patients treated with cyclosporine. The goal of therapy should be LDL-C levels less than 100 mg/dL. However, if the LDL-C levels remain 130 mg/dL or greater, consider adding a second agent to the statin (eg, a bile acid sequestrant¹³ or

Figure 5

National Kidney Foundation Treatment Guidelines for Adult Renal Transplant Recipients With Dyslipidemias¹³



TG = triglycerides; TLC = therapeutic lifestyle changes; LDL-C = low-density lipoprotein cholesterol; HDL-C = high-density lipoprotein cholesterol.

Adapted with permission from K/DOQI clinical practice guidelines for management of dyslipidemias in patients with kidney disease. *Am J Kidney Dis.* 2003;41:S1-S91.

ezetimibe). In a preliminary study of 40 stable kidney transplant recipients, ezetimibe treatment did lower LDL-C levels⁵⁶; however, ezetimibe use with cyclosporine is not recommended for cardiac transplant recipients because of a pharmacokinetic interaction that leads to a 12-fold increase in ezetimibe levels.^{57,58} For some patients who continue to have high LDL-C levels despite therapy, especially if the risk for ischemic heart disease appears to outweigh the risk of acute rejection, consideration may be given to changing immunosuppression.

The role of immunosuppressive drugs in dyslipidemias

Corticosteroids, cyclosporine, and sirolimus can each contribute to dyslipidemias after kidney transplantation. However, these and other immunosuppressive agents can also adversely affect other cardiovascular disease risk factors (Table 1, page 3). Of course, the risk for rejection and graft dysfunction or failure must also be considered in selecting immunosuppressive agents. In the end, the clinician is left with a delicate balancing act of choosing between different risks and benefits from various immunosuppressive drug combinations.

Corticosteroids

Corticosteroids have long been associated with increased lipid levels, making immunosuppressive medication protocols that minimize the use of prednisone attractive. In a multicenter, randomized 6-month study of controlled corticosteroid dose reduction in kidney transplant recipients also receiving MMF and cyclosporine, a significant reduction in total cholesterol levels was observed in the group who had a 50% corticosteroid dose reduction then cessation (low/stop group, dose of 0-5 mg/day) compared to the control group (dose of 10 mg/day) who continued to receive corticosteroids (Figure 6). Systolic BP was also lower in the low/stop group, and there was a reduction in other corticosteroid-related side effects, such as bone loss. Although at 6 months, the low/stop group experienced a significantly higher rate of acute rejection than did the control group (23% vs 14%, respectively; $P=.008$), the rejections were mostly Banff grade I.⁵⁹

In another multicenter study from Europe, investigators compared lipid levels at 3 and 6 months posttransplantation in a tacrolimus-based regimen that withdrew corticosteroids or MMF after 3 months. Immunologically low-risk patients were followed for an additional 3 months. Reductions in mean total cholesterol and LDL-C levels were significantly greater in the corticosteroid-withdrawal group (MMF + tacrolimus) than in either the control


(corticosteroids + MMF + tacrolimus) or the MMF-withdrawal (corticosteroids + tacrolimus) group ($P<.001$). However, the changes in mean HDL-C levels resulted in little final change in the LDL-C/HDL-C ratio in the three groups between 4 and 6 months.³²

Calcineurin and target of rapamycin inhibitors

There is now substantial evidence that the CNIs tacrolimus, unlike cyclosporine, does not affect lipid levels. This difference was demonstrated in a multicenter study in which stable kidney transplant recipients with established hyperlipidemia were randomized for conversion from cyclosporine-based immunosuppression to a tacrolimus-based regimen or continued treatment on the cyclosporine-based regimen. After 6 months of follow-up in patients converted to tacrolimus ($n=27$), total cholesterol levels decreased by 55 mg/dL (-16%, $P=.003$) and LDL-C levels decreased by 48 mg/dL (-25%, $P=.001$) compared to levels in recipients remaining on cyclosporine-based immunosuppression ($n=26$).⁶⁰ This finding has also been reported by other groups.⁶¹⁻⁶³ However, these results should be cautiously interpreted pending further understanding of the risk factors for cardiovascular disease in the transplant population.⁶⁴

The target of rapamycin inhibitor sirolimus can cause elevations in lipoproteins that may be marked in some cases.⁶⁵ One randomized trial comparing sirolimus- and cyclosporine-based regimens reported that both groups showed significant increases in total cholesterol, HDL-C, and LDL-C levels by 1 month posttransplantation, compared with pretransplant levels ($P=.001$). Additionally, 64.5% of sirolimus-treated patients and 53.3% of cyclosporine-treated patients

Figure 6
Cholesterol Levels After Prednisone Withdrawal in Cyclosporine/Mycophenolate Mofetil-Treated Renal Transplant Recipients⁵⁹



received lipid-lowering drug therapy within 1 year post-transplantation.⁶⁶ Another trial randomized recipients to tacrolimus-based immunosuppression (and corticosteroids) with either MMF or sirolimus for 6 months. Elevated concentrations of total cholesterol ($P=.0001$) and LDL-C ($P=.001$) were significantly more prevalent in the sirolimus-treated patients than in the MMF-treated group.³⁰ However, a number of positive effects of sirolimus, in particular on graft function, make its use a viable option for many patients.⁶⁷

Role of lipid-lowering drugs

A substantial amount of data from randomized controlled trials in the general population indicates that statins are very effective in reducing levels of total cholesterol and LDL-C, and cardiovascular disease events. However, there is some uncertainty about the pathogenesis of cardiovascular disease in transplant recipients, or in those patients with chronic kidney disease, and whether treatment with statins will have similar efficacy to that in the general population. Results from “Die deutsche Diabetes Dialyse Studie” (4D) trial in patients with diabetes and ESRD showed no significant difference in the composite endpoint of major adverse cardiac events (MACE) in patients receiving atorvastatin compared to placebo.^{68,69} The Assessment of Lescol in Renal Transplantation randomized controlled trial investigated the effect of fluvastatin on cardiac outcomes in kidney transplant recipients. This adequately powered study followed 2102 clinically stable transplant recipients (with total cholesterol levels between 155 mg/dL and 348 mg/dL) receiving fluvastatin (40 mg daily) or placebo for a mean follow-up of 5.1 years. Although fluvastatin lowered LDL-C levels by 32% and reduced the incidence of some cardiac events, the risk reduction for the primary endpoint of MACE was not significantly different from that in the placebo group (Table 4).⁷⁰ A further analysis failed to show that fluvastatin preserved graft function.^{71,72}

Another critical outcome measure in kidney transplantation is the incidence of acute rejection. A number of studies investigated the possible effect of statin therapy on acute rejection. Only one study, with an unusually high rate of rejection in the control group, showed a significant improvement in acute rejection rates with statin therapy ($P=.01$),⁷³ whereas a number of recent studies have found no difference.^{74,75} Thus, it does not appear that statins reduce the incidence of acute rejection in kidney transplant recipients.

The pharmacokinetic interaction between CNIs and statins, documented for cyclosporine and possibly tacrolimus, results in a higher blood level of the statin.^{13,76} As a result, reduced doses of statins should probably be used in patients also treated with cyclosporine. Additionally, the fact that both statins and CNIs are metabolized by the CYP450 pathway can result in unwanted interactions with other drugs metabolized by the same pathway, such as CCBs and azole antifungals.¹³

In instances when treatment with diet and a statin does not achieve the goal of LDL-C levels less than 100 mg/dL, consideration should be given to adding a second agent. A bile acid sequestrant can be used effectively with a statin, but bile acid sequestrants are often not well tolerated.¹³ The new cholesterol uptake inhibitor ezetimibe may be used in combination with a statin. Levels of ezetimibe may be increased in cyclosporine-treated patients, but the consequences of this are unknown.^{56,57} Fibrates lower LDL-C levels in the general population, but increased blood levels of statins in patients also receiving cyclosporine and a statin make adding a fibrate to a statin in cyclosporine-treated patients risky.¹³

CONCLUSIONS

Cardiovascular disease is a prevalent, life-threatening issue for kidney transplant recipients. Neither hypertension nor dyslipidemias are well controlled. The immunosuppressive drugs critical for survival of the allograft cause both hypertension and dyslipidemias in kidney transplant recipients. TLC, together with BP- and lipid-lowering agents, are needed in the majority of transplant recipients. In the future, effective immunosuppressive agents that can replace corticosteroids and CNIs may help reduce cardiovascular disease risk.

Table 4 Endpoints in the ALERT Trial of Fluvastatin Therapy in Renal Transplant Recipients⁷⁰

Endpoint	P	Risk Ratio (95% CI)
Cardiac death	.031	0.62 (0.40-0.96)
Definite nonfatal MI	.050	0.68 (0.40-1.00)
Cardiac death or definite nonfatal MI	.005	0.65 (0.48-0.88)
Cardiac death or definite or probable nonfatal MI	.032	0.72 (0.54-0.97)
CABG	.932	1.03 (0.58-1.81)
PCI	.357	0.80 (0.49-1.30)
Cardiac death or definite or probable nonfatal MI, CABG/PCI (MACE)	.139	0.83 (0.64-1.06)
Fatal or nonfatal cerebrovascular events*	.391	1.16 (0.83-1.63)
Noncardiovascular deaths	.288	1.20 (0.86-1.67)
All-cause death	.848	1.02 (0.81-1.30)
Graft loss or doubling of serum creatinine	.369	1.10 (0.89-1.36)

ALERT = Assessment of Lescol in Renal Transplantation; CI = confidence interval; MI = myocardial infarction; CABG = coronary artery bypass grafting; PCI = percutaneous coronary intervention; MACE = major adverse coronary events. *Fatal or nonfatal stroke, transient ischemic attack, reversible ischemic neurologic deficit, subarachnoid hemorrhage. Adapted with permission from Holdaas H, et al. *Lancet*. 2003;361:2024-2031.

REFERENCES

- Baroletti SA, Gabardi S, Magee CC, Milford EL. Calcium channel blockers as the treatment of choice for hypertension in renal transplant recipients: fact or fiction. *Pharmacotherapy*. 2003;23:788-801.
- Wolfe RA, Ashby VB, Milford EL, et al. Comparison of mortality in all patients on dialysis, patients on dialysis awaiting transplantation, and recipients of a first cadaveric transplant. *N Engl J Med*. 1999;341:1725-1730.
- Foley RN, Parfrey PS, Sarnak MJ. Clinical epidemiology of cardiovascular disease in chronic renal disease. *Am J Kidney Dis*. 1998;32(5 suppl 3):S112-S119.
- Foley RN, Parfrey PS, Sarnak MJ. Epidemiology of cardiovascular disease in chronic renal disease. *J Am Soc Nephrol*. 1998;9(12 suppl):S16-S23.
- Kasiske BL, Chakkera HA, Roel J. Explained and unexplained ischemic heart disease risk after renal transplantation. *J Am Soc Nephrol*. 2000;11:1735-1743.
- Castelli WP. Epidemiology of coronary heart disease: the Framingham study. *Am J Med*. 1984;76:4-12.
- Curtis JJ. Hypertensinogenic mechanism of the calcineurin inhibitors. *Curr Hypertens Rep*. 2002;4:377-380.
- Midtvedt K, Neumayer HH. Management strategies for posttransplant hypertension. *Transplantation*. 2000;70(11 suppl):SS64-SS69.
- Meier-Kriesche HU, Baliga R, Kaplan B. Decreased renal function is a strong risk factor for cardiovascular death after renal transplantation. *Transplantation*. 2003;75:1291-1295.
- Zhang R, Leslie B, Boudreaux JP, Frey D, Reisn E. Hypertension after kidney transplantation: impact, pathogenesis and therapy. *Am J Med Sci*. 2003;325:202-208.
- K/DOQI clinical practice guidelines on hypertension and antihypertensive agents in chronic kidney disease. Available at: http://www.kidney.org/professionals/kdoqi/guidelines_bp/. Accessed March 16, 2005.
- K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis*. 2002;39(2 suppl 1):S1-S266.
- K/DOQI clinical practice guidelines for management of dyslipidemias in patients with kidney disease. *Am J Kidney Dis*. 2003;41(4 suppl 3):1-IV, S1-S91.
- Implications of preserving long-term renal function after renal transplantation: epidemiology of renal dysfunction and cardiovascular disease. Available at: <http://www.medscape.com/viewprogram/3016>. Accessed March 16, 2005.
- Kasiske BL, Anjum S, Shah R, et al. Hypertension after kidney transplantation. *Am J Kidney Dis*. 2004;43:1071-1081.
- Haydar AA, Covic A, Jayawardene S, et al. Insights from ambulatory blood pressure monitoring: diagnosis of hypertension and diurnal blood pressure in renal transplant recipients. *Transplantation*. 2004;77:849-853.
- Opelz G, Wujciak T, Ritz E. Association of chronic kidney graft failure with recipient blood pressure. Collaborative Transplant Study. *Kidney Int*. 1998;53:217-222.
- Mange KC, Cizman B, Joffe M, Feldman HI. Arterial hypertension and renal allograft survival. *JAMA*. 2000;283:633-638.
- Zeier M, Mandelbaum A, Ritz E. Hypertension in the transplanted patient. *Nephron*. 1998;80:257-268.
- Curtis JJ, Lucas BA, Kotchen TA, Luke RG. Surgical therapy for persistent hypertension after renal transplantation. *Transplantation*. 1981;31:125-128.
- Bruno S, Remuzzi G, Ruggenenti P. Transplant renal artery stenosis. *J Am Soc Nephrol*. 2004;15:134-141.
- Textor SC, Taler SJ, Driscoll N, et al. Blood pressure and renal function after kidney donation from hypertensive living donors. *Transplantation*. 2004;78:276-282.
- Guyton AC. Abnormal renal function and autoregulation in essential hypertension. *Hypertension*. 1991;18(5 suppl):III49-III53.
- Curtis JJ, Luke RG, Dustan HP, et al. Remission of essential hypertension after renal transplantation. *N Engl J Med*. 1983;309:1009-1015.
- Veenstra DL, Best JH, Hornberger J, Sullivan SD, Hricak DE. Incidence and long-term cost of steroid-related side effects after renal transplantation. *Am J Kidney Dis*. 1999;33:829-839.
- Nichelle L, Canet S, Garrigue V, Chong G, Mourad G. Arterial hypertension in renal transplant recipients treated with tacrolimus or cyclosporine-Neoral. *Transplant Proc*. 2002;34:2824-2825.
- Zhang W, Victor RG. Calcineurin inhibitors cause renal afferent activation in rats: a novel mechanism of cyclosporine-induced hypertension. *Am J Hypertens*. 2000;13:999-1004.
- Halloran PF. Immunosuppressive drugs for kidney transplantation. *N Engl J Med*. 2004;351:2715-2729.
- Margreiter R, for the European Tacrolimus vs Cyclosporin Microemulsion Renal Transplantation Study Group. Efficacy and safety of tacrolimus compared with cyclosporin microemulsion in renal transplantation: a randomised multicentre study. *Lancet*. 2002;359:741-746.
- Gonwa T, Mendez R, Yang HC, Weinstein S, Jensik S, Steinberg S, for the Prograf Study Group. Randomized trial of tacrolimus in combination with sirolimus or mycophenolate mofetil in kidney transplantation: results at 6 months. *Transplantation*. 2003;75:1213-1220.
- Oberbauer R, Kreis H, Johnson RW, et al. Long-term improvement in renal function with sirolimus after early cyclosporine withdrawal in renal transplant recipients: 2-year results of the Rapamune Maintenance Regimen Study. *Transplantation*. 2003;76:364-370.
- Vanrenterghem Y, van Hooff JP, Squifflet JP, et al. Minimization of immunosuppressive therapy after renal transplantation: results of a randomized controlled trial. *Am J Transplant*. 2005;5:87-95.
- K/DOQI clinical practice guidelines on hypertension and antihypertensive agents in chronic kidney disease. Available at: http://www.kidney.org/professionals/kdoqi/guidelines_bp/. Accessed March 16, 2005.
- European best practice guidelines for renal transplantation. Section IV: long-term management of the transplant recipient. IV.5.2. Cardiovascular risks. Arterial hypertension. *Nephrol Dial Transplant*. 2002;17(suppl 4):25-26.
- Curtis JJ, Luke RG, Whelchel JD, Diethelm AG, Jones P, Dustan HP. Inhibition of angiotensin-converting enzyme in renal-transplant recipients with hypertension. *N Engl J Med*. 1983;308:377-381.
- Formica RN Jr, Friedman AL, Lorber MI, Bia MJ. Angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers used for the treatment of hypertension appear to be safe in the early posttransplant period. *Transplant Proc*. 2004;36:2675-2678.
- Stigant CE, Cohen J, Vivera M, Zaltzman JS. ACE inhibitors and angiotensin II antagonists in renal transplantation: an analysis of safety and efficacy. *Am J Kidney Dis*. 2000;35:58-63.
- Grekas D, Dioudis C, Kalevosoglou I, Alivannis P, Derveniotis V, Tourkantonis A. Renal hemodynamics in hypertensive renal allograft recipients: effects of calcium antagonists and ACE inhibitors. *Kidney Int Suppl*. 1996;55:S97-S100.
- Premasathian NC, Muehrer R, Brazzy PC, Pirsch JD, Becker BN. Blood pressure control in kidney transplantation: therapeutic implications. *J Hum Hypertens*. 2004;18:871-877.
- Reil K, Linde J, Morzycka-Michalik M, Gaciong Z, Lao M. Effect of enalapril on proteinuria after kidney transplantation. *Transpl Int*. 1993;6:213-217.
- el-Agroud AE, Hassan NA, Foda MA, et al. Effect of angiotensin II receptor blocker on plasma levels of TGF-beta 1 and interstitial fibrosis in hypertensive kidney transplant patients. *Am J Nephrol*. 2003;23:300-306.
- Rustom R, Grime JS, Sells RA, et al. Renal tubular peptide catabolism in chronic vascular rejection. *Ren Fail*. 2001;23:517-531.
- Nigbor DA, Lewis JB. Use of calcium antagonists in renal patients: therapeutic benefit or medical malpractice? *Curr Hypertens Rep*. 2003;5:430-436.
- van Riemsdijk IC, Mulder PG, de Fijter JW, et al. Addition of isradipine (Lomir) results in a better renal function after kidney transplantation: a double-blind, randomized, placebo-controlled, multi-center study. *Transplantation*. 2000;70:122-126.
- Chanard J, Toupance O, Lavaud S, Hurault de Ligny B, Bernaud C, Moulin B. Amlodipine reduces cyclosporin-induced hyperuricaemia in hypertensive renal transplant recipients. *Nephrol Dial Transplant*. 2003;18:2147-2153.
- Harper SJ, Moorhouse J, Veitch PS, et al. Nifedipine improves immediate, and 6- and 12-month graft function in cyclosporin A (CyA) treated renal allograft recipients. *Transpl Int*. 1992;5(suppl 1):S69-S72.
- Nyberg G, Haljamae U, Herlitz H, Norden G, Blohme I. Early renal graft function in recipients treated with the calcium channel blocker felodipine. *Scand J Urol Nephrol*. 1994;28:17-20.
- Sica DA, Douglas JG. The African American Study of Kidney Disease and Hypertension (AASK): new findings. *J Clin Hypertens (Greenwich)*. 2001;3:244-251.
- Midtvedt K, Hartmann A, Foss A, et al. Sustained improvement of renal graft function for two years in hypertensive renal transplant recipients treated with nifedipine as compared to lisinopril. *Transplantation*. 2001;72:1787-1792.
- Moore R, Thomas D, Morgan E, et al. Abnormal lipid and lipoprotein profiles following renal transplantation. *Transplant Proc*. 1993;25(1 pt 2):1060-1061.
- Brown JH, Murphy BG, Douglas AF, et al. Influence of immunosuppressive therapy on lipoprotein(a) and other lipoproteins following renal transplantation. *Nephron*. 1997;75:277-282.
- Roodnat JL, Mulder PG, Zietse R, et al. Cholesterol as an independent predictor of outcome after renal transplantation. *Transplantation*. 2000;69:1704-1710.
- Massy ZA, Gujjarro C, Wiederkehr MR, Ma JZ, Kasiske BL. Chronic renal allograft rejection: immunologic and nonimmunologic risk factors. *Kidney Int*. 1996;49:518-524.
- Stamler J, Wentworth D, Neaton JD. Is relationship between serum cholesterol and risk of premature death from coronary heart disease continuous and graded? Findings in 356,222 primary screenees of the Multiple Risk Factor Intervention Trial (MRFIT). *JAMA*. 1986;256:2823-2828.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA*. 2001;285:2486.
- Puthenparumpil JJ, Keough-Ryan T, Kiberd M, Lawen J, Kiberd BA. Treatment of hypercholesterolemia with ezetimibe in the kidney transplant population. *Transplant Proc*. 2005;37:1033-1035.
- Bilchick KC, Henrikson CA, Skojec D, Kasper EK, Blumenthal RS. Treatment of hyperlipidemia in cardiac transplant recipients. *Am Heart J*. 2004;148:200-210.
- Ezetimibe prescribing information. Available at: http://www.zetia.com/ezetimibe/zetia/consumer/product_information/pi/index.jsp. Accessed March 16, 2005.
- Vanrenterghem Y, Lebranchu Y, Hene R, Oppenheimer F, Ekberg H. Double-blind comparison of two corticosteroid regimens plus mycophenolate mofetil and cyclosporine for prevention of acute renal allograft rejection. *Transplantation*. 2000;70:1352-1359.
- McCune TR, Thacker LR II, Peters TG, et al. Effects of tacrolimus on hyperlipidemia after successful renal transplantation: a Southeastern Organ Procurement Foundation multicenter clinical study. *Transplantation*. 1998;65:87-92.
- Baid-Agrawal S, Delmonico FL, Tolkoff-Rubin NE, et al. Cardiovascular risk profile after conversion from cyclosporine A to tacrolimus in stable renal transplant recipients. *Transplantation*. 2004;77:1199-1202.
- Raofi V, Holman DM, Coady N, et al. A prospective randomized trial comparing the efficacy of tacrolimus versus cyclosporine in black recipients of primary cadaveric renal transplants. *Am J Surg*. 1999;177:299-302.
- Artz MA, Boots JM, Ligtienberg G, et al. Improved cardiovascular risk profile and renal function in renal transplant patients after randomized conversion from cyclosporine to tacrolimus. *J Am Soc Nephrol*. 2003;14:1880-1888.
- Cosio FG, Larson TS. Cardiovascular disease after transplantation: do we know all of the variables? *Am J Transplant*. 2003;3:915-916.
- Kreis H, Cisterne JM, Land W, et al. Sirolimus in association with mycophenolate mofetil induction for the prevention of acute graft rejection in renal allograft recipients. *Transplantation*. 2000;69:1252-1260.
- Flechner SM, Goldfarb D, Modlin C, et al. Kidney transplantation without calcineurin inhibitor drugs: a prospective, randomized trial of sirolimus versus cyclosporine. *Transplantation*. 2002;74:1070-1076.
- Morales JM, Wranner L, Kreis H, et al. Sirolimus does not exhibit nephrotoxicity compared to cyclosporine in renal transplant recipients. *Am J Transplant*. 2002;2:436-442.
- Wanner C, Krane V, Marz W, et al. for the Deutsche Diabetes-Dialyse-Studie (4D) Study Group. Randomized controlled trial on the efficacy and safety of atorvastatin in patients with type 2 diabetes on hemodialysis (4D study): demographic and baseline characteristics. *Kidney Blood Press Res*. 2004;27:259-266.
- Atorvastatin lowers LDL but not risk of CV events in diabetics with ESRD. Available at: <http://www.lifescan.com/professionals/hcp/news/dn110104-1/>. Accessed March 16, 2005.
- Holdaas H, Fellström B, Jardine AG, et al. Effect of fluvastatin on cardiac outcomes in renal transplant recipients: a multicentre, randomised, placebo-controlled trial. *Lancet*. 2003;361:2024-2031.
- Fellström B, Holdaas H, Jardine AG, et al. Effect of fluvastatin on renal endpoints in the Assessment of Lescol in Renal Transplantation (ALERT) trial. *Kidney Int*. 2004;66:1549-1555.
- Fellström B, Holdaas H, Jardine AG, et al. Risk factors for reaching renal endpoints in the Assessment of Lescol in Renal Transplantation (ALERT) trial. *Transplantation*. 2005;79:205-212.
- Katznelson S, Wilkinson AH, Kobashigawa JA, et al. The effect of pravastatin on acute rejection after kidney transplantation—a pilot study. *Transplantation*. 1996;61:1469-1474.
- Kasiske BL, Heim-Duthoy KL, Singer GG, Watschinger B, Germain MJ, Bastani B. The effects of lipid-lowering agents on acute renal allograft rejection. *Transplantation*. 2001;72:223-227.
- Sahu K, Sharma R, Gupta A, et al. Effect of lovastatin, an HMG CoA reductase inhibitor, on acute renal allograft rejection. *Clin Transplant*. 2001;15:173-175.
- Asberg A. Interactions between cyclosporin and lipid-lowering drugs: implications for organ transplant recipients. *Drugs*. 2003;63:367-378.

STATE-OF-THE-ART MANAGEMENT OF POSTTRANSPLANT SEQUELAE

IMPACT AND MANAGEMENT OF CARDIOVASCULAR RISKS AFTER KIDNEY TRANSPLANTATION

CME/CE POSTTEST AND EVALUATION

Release Date: June 2005 Expiration Date: June 30, 2006

If you wish to receive CME/CE credit and a statement of completion, please mail or fax a copy of your completed answer sheet/registration/evaluation on page 10 to:

For physicians and nurses: University of Minnesota
Office of CME
200 Oak Street SE, Suite 190
Minneapolis, MN 55455
Attn: Distance Learning (DL-05-105B)
Fax: 612-626-7766

For pharmacists: Continuing Pharmacy Education
University of Minnesota
College of Pharmacy
420 Delaware Street SE, MMC 387
Minneapolis, MN 55455
Fax: 612-626-4613

POSTTEST

Rebecca L. is a 35-year-old kidney transplant recipient (height, 5 foot 4 in., weighs 182 lb.) who has been engaged for 4 months. In part because of the sedentary nature and long work hours of her job as an auditor, she has gained 25 lbs in the 2.5 years since receiving her transplant. She has begun smoking again to try to curb her appetite. Rebecca lost her native kidneys from idiopathic nephrotic syndrome and was fortunate to have a cousin donate a well-matched kidney for transplantation. Rebecca talked to her nephrologist during a scheduled check-up about approaches to help her lose weight before her wedding. Her posttransplant clinical course included one episode of mild acute rejection (Banff I) within 3 months posttransplantation, which was resolved with pulse methylprednisone, and she has otherwise

had an uneventful history beyond weight gain. Rebecca is currently on a maintenance immunosuppressive regimen of cyclosporine 3 to 3.5 mg/day (adjusted to trough whole blood levels of 150 to 200 ng/mL), mycophenolate mofetil (1 mg bid), and prednisone (7.5 mg/day). Blood tests taken at her recent appointment revealed that her total cholesterol (from 200 to 230 mg/dL), low-density lipoprotein cholesterol (from 100 to 130 mg/dL), and triglyceride (from 160 to 180 mg/dL) levels have gradually increased over the last 12 months. Blood pressure is under control (130/76 mm Hg) with an angiotensin-converting enzyme (ACE) inhibitor. Renal function remains good.

- Which nonimmunosuppression-related factors should be discussed with Rebecca?
 - Nutrition counseling
 - Smoking cessation
 - Increasing exercise
 - All of the above
 - a and b
- If cholesterol levels are not reduced by therapeutic lifestyle counseling, what would be your next approach?
 - Add statin therapy
 - Add fibrate therapy
 - Withdraw mycophenolate mofetil
 - Minimize corticosteroids
 - Switch from the ACE inhibitor to a calcium channel blocker
- Renal transplant recipients have:
 - A cardiovascular mortality rate of approximately 40%
 - An overall higher cardiovascular mortality rate than the general population
 - A nontraditional risk factor for cardiovascular disease related to some immunosuppressive drugs
 - None of the above
 - All of the above
- Hypertension in renal transplant recipients:
 - Is an independent risk factor for poorer long-term kidney graft survival
 - Is frequently underdetected through routine clinic visit measurements of blood pressure compared to ambulatory blood pressure monitoring
 - Can exhibit blood pressure changes with a diurnal variation similar to that in patients with chronic kidney disease
 - None of the above
 - All of the above
- Use of calcineurin inhibitors (CNIs) in renal transplantation resulted in:
 - Continued CNI maintenance therapy in all patients without drug minimization opportunities
 - No changes in blood pressure compared to pretransplant values
 - Studies showing that tacrolimus induced significantly more hypertension than did cyclosporine
 - None of the above
 - All of the above
- Target blood pressure for kidney transplant recipients recommended in the National Kidney Foundation guidelines is:
 - <145/90 mm Hg
 - <140/85 mm Hg
 - <140/80 mm Hg
 - <135/85 mm Hg
 - <130/80 mm Hg
- Since Rebecca's lipid profile deteriorated during the last 12 months, would you modify her immunosuppressive protocol?
 - Yes, eliminate cyclosporine
 - Yes, eliminate sirolimus
 - Yes, decrease cyclosporine doses
 - Yes, decrease sirolimus doses
 - Would not change immunosuppressive protocol
- In renal transplant recipients:
 - Studies have shown little benefit for calcium channel blockers in comparison to diuretics, angiotensin-converting enzyme (ACE) inhibitors, or angiotensin II receptor blockers for antihypertensive therapy
 - ACE inhibitor use can reduce proteinuria
 - Diuretics should not be used in cases of volume overload and sodium retention
 - ACE inhibitors increase hematocrit or hemoglobin levels in those patients with posttransplant erythrocytosis
 - Only 5% of patients will need to use more than one antihypertensive drug to manage blood pressure
- Cholesterol levels were reduced by minimizing or eliminating the following drug(s) in immunosuppressive regimens after kidney transplantation:
 - Cyclosporine
 - Corticosteroids
 - Sirolimus
 - None of the above
 - All of the above
- Use of lipid-lowering therapies, immunosuppressive drugs, and other commonly used drugs must be approached carefully as undesirable pharmacokinetic interactions are known to occur between:
 - Cyclosporine and high-dose statins
 - Fibrates and statins and cyclosporine
 - Statins and cyclosporine and azole antifungals
 - All of the above
 - None of the above

STATE-OF-THE-ART MANAGEMENT OF POSTTRANSPLANT SEQUELAE

IMPACT AND MANAGEMENT OF CARDIOVASCULAR RISKS AFTER KIDNEY TRANSPLANTATION

CME/CE POSTTEST AND EVALUATION

Release Date: June 2005 Expiration Date: June 30, 2006

POSTTEST ANSWER KEY (questions from page 9)

- | | | | | |
|--------------|--------------|--------------|--------------|---------------|
| 1. a b c d e | 3. a b c d e | 5. a b c d e | 7. a b c d e | 9. a b c d e |
| 2. a b c d e | 4. a b c d e | 6. a b c d e | 8. a b c d e | 10. a b c d e |

PROGRAM EVALUATION

The University of Minnesota would appreciate your comments regarding the quality of the information presented.

1. Each of the following program's educational objectives were fully met:

- Recognize the prevalence and long-term impact of overall cardiovascular disease, and specifically hypertension and dyslipidemias, in renal transplant recipients
☐ Strongly Agree ☐ Agree ☐ Disagree ☐ Strongly Disagree
- Describe the risk factors for hypertension and dyslipidemias
☐ Strongly Agree ☐ Agree ☐ Disagree ☐ Strongly Disagree
- Identify the effects of immunosuppressive drugs on the development of hypertension and dyslipidemias
☐ Strongly Agree ☐ Agree ☐ Disagree ☐ Strongly Disagree
- Outline the modifications to immunosuppressive regimens to help manage hypertension and dyslipidemias
☐ Strongly Agree ☐ Agree ☐ Disagree ☐ Strongly Disagree
- Discuss the interventions to treat hypertension and dyslipidemias
☐ Strongly Agree ☐ Agree ☐ Disagree ☐ Strongly Disagree

2. The quality of the educational process (method of presentation and information provided) was satisfactory and appropriate.

☐ Strongly Agree ☐ Agree ☐ Disagree ☐ Strongly Disagree

3. The educational activity has enhanced my professional effectiveness and improved my ability to treat/manage patients.

☐ Strongly Agree ☐ Agree ☐ Disagree ☐ Strongly Disagree ☐ N/A

4. The educational activity has improved my ability to communicate with patients.

☐ Strongly Agree ☐ Agree ☐ Disagree ☐ Strongly Disagree ☐ N/A

5. The information presented was free of promotional or commercial bias.

☐ Agree ☐ Disagree

6. What changes will you make in your practice as a result of participating in this program?

7. Comments/suggestions regarding *this* material: _____

8. Recommendations for *future* presentations: _____

9. What is the most important barrier to the optimal posttransplant management of patients receiving renal transplants? (select one answer)

- | | |
|---|-------------------------------------|
| <input type="checkbox"/> Patient adherence | <input type="checkbox"/> Infections |
| <input type="checkbox"/> Side effects of immunosuppressive agents | <input type="checkbox"/> Neoplasia |
| <input type="checkbox"/> Renal function | <input type="checkbox"/> Other |

10. Approximately how many patients do you see per week? _____

11. Approximately what percentage of your patients are renal transplant recipients?

Degree: ☐ MD ☐ RN ☐ Other

☐ DO ☐ PharmD _____

Job Title

- ☐ Transplant Surgeon
☐ Nephrologist
☐ Transplant Coordinator
☐ Transplant Pharmacist
☐ Transplant Case Manager
☐ Nurse
☐ Other _____

Practice Type

- ☐ Private
☐ Group
☐ Transplant Center
☐ Academic

Practice Location

- ☐ Urban
☐ Suburban
☐ Rural

Full Name _____

Company/Affiliation _____

Street Address _____

City _____

State _____

ZIP Code _____

Email Address _____

Fax Number _____

I certify that I completed this CME/CE activity. The actual amount of time I spent in this activity was: _____ hours _____ minutes.

Signature _____

Date Completed _____

PHYSICIANS: Are you licensed in the United States?

☐ YES

☐ NO

NURSES: State of license and number _____

The information presented in this material is intended solely for the continuing medical education needs of healthcare professionals. Healthcare professionals and individuals should not rely upon any of the information provided in this material. Some presented product information may be for unlabeled/investigational uses. Before using or prescribing any product discussed in this publication, clinicians should consult the full prescribing information.

The views presented herein are those of the faculty and not necessarily those of SynerMed® Communications, the commercial supporter, or CME sponsor.

SynerMed® Communications owns all copyrights in this material, and no person shall have the right to use, duplicate, distribute, modify, or create derivative versions of this material, in any manner or in any medium, except as necessary to complete the program for obtaining continuing medical education credit. Any violation of this shall result in appropriate legal action being taken.

**All correspondence concerning the contents of
this publication should be directed to:**

**SynerMed® Communications
Dept. WA48D
405 Trimmer Road
PO Box 458
Califon, NJ 07830**



Developed and Produced by

